

The SIDDHARTA-2 experiment: preparation for the first kaonic deuterium measurement

L. De Paolis,^{a,*} M. Bazzi,^a D. Bosnar,^b M. Bragadireanu,^c M. Cargnelli,^d M. Carminati,^e A. Clozza,^a G. Deda,^e R. Del Grande,^{f,a} K. Dulski,^a C. Fiorini,^e I. Friščić,^b C. Guaraldo,^a M. Iliescu,^a M. Iwasaki,^g P. King,^e A. Khreptak,^{a,i} P. Levi Sandri,^a S. Manti,^a J. Marton,^d M. Miliucci,^a P. Moskal,^{h,i} F. Napolitano,^a S. Niedźwiecki,^{h,i} H. Onishi,^j K. Piscicchia,^k Y. Sada,^j A. Scordo,^a F. Sgaramella,^a H. Shi,^d M. Silarski,ⁱ D. L. Sirghi,^{k,a} F. Sirghi,^a M. Skurzok,^{h,i} A. Spallone,^a K. Toho,^g M. Tüchler,^{d,l} C. Yoshida,^j J. Zmeskal^d and C. Curceanu^a

^aINFN, Laboratori Nazionali di Frascati, Via E. Fermi 54, I-00044 Frascati(RM), Italy

^bDepartment of Physics, Faculty of Science, University of Zagreb, 10000 Zagreb, Croatia

^cHoria Hulubei National Institute of Physics and Nuclear Engineering, Str. Atomistilor No. 407, P.O. Box MG-6 Bucharest-Măgurele, Romania

^dStefan Meyer Institute for Subatomic Physics, Kegelgasse 27, 1030 Wien, Austria

^ePolitecnico di Milano, Dipartimento di Elettronica, Informazione e Bioingegneria and INFN Sezione di Milano, I-20133 Milano, Italy

^fPhysik Department E62, Technische Universität München, James-Frank-straße 1, 85748 Garching, Germany

^gRIKEN, Institute of Physical and Chemical Research, Wako, Tokyo 351-0198, Japan

^hCenter for Theranostics, Jagiellonian University, Krakow, Poland

ⁱFaculty of Physics, Astronomy, and Applied Computer Science, Jagiellonian University, Krakow, 30-348, Poland

^jResearch Center for Electron Photon Science (ELPH), Tohoku University, Sendai, Japan

^kCentro Ricerche Enrico Fermi—Museo Storico della Fisica e Centro Studi e Ricerche “Enrico Fermi”, Via Panisperna 89a, I-00184 Roma, Italy

^lUniversity of Vienna, Vienna Doctoral School in Physics, Vienna, Austria

E-mail: Luca.DePaolis@lnf.infn.it

The SIDDHARTA-2 experiment aims to perform the first measurement of the kaonic deuterium $2p \rightarrow 1s$ x-ray transitions. The apparatus is presently installed at the interaction region of the DAΦNE electron-positron collider at the National Laboratories of Frascati (LNF-INFN), in Italy. Kaonic deuterium and kaonic hydrogen $2p \rightarrow 1s$ x-ray transitions measurements, the latter one already performed by the SIDDHARTA collaboration, allow the determination of antikaon-nucleon scattering lengths. A description of the SIDDHARTA-2 apparatus in preparation for the kaonic deuterium measurement is provided in this paper.

41st International Conference on High Energy physics - ICHEP2022

6-13 July, 2022

Bologna, Italy

*Speaker

1. Introduction

The x-ray spectroscopy of light kaonic atoms allows the experimental investigation of the strong interaction between the kaon and the nucleus at low energies [1]. A kaonic atom is formed when a K^- is stopped in a dedicated target and captured in an atomic system due to the electromagnetic interaction with the nucleus. The captured K^- replaces an electron in a highly excited atomic level and initiates an electromagnetic cascade process that could bring it to the innermost atomic level, in which the kaon-nucleus strong interaction produces a shift and a broadening of the energy level that can be extracted with dedicated x-ray spectroscopy measurements. In 2009, the SIDDHARTA experiment measured the kaonic hydrogen $2p \rightarrow 1s$ transition, extracting the energy shift and width of the $1s$ kaonic hydrogen atomic level due to the strong kaon-proton interaction [2]. This measurement, added to the kaonic deuterium $2p \rightarrow 1s$ transition measurement, will allow to extract the antikaon-nucleon scattering lengths, not directly accessible otherwise due to the $\Lambda(1405)$ resonance just below the threshold and the strong coupling to the $\pi\Sigma$ channel [1, 3]. The SIDDHARTA-2 experiment aims to measure, for the first time, the kaonic deuterium $2p \rightarrow 1s$ transition in 2023/4 at the DAΦNE electron-positron collider [4], a Φ factory located at the National Laboratories of Frascati, in Italy. The SIDDHARTA-2 apparatus, presently installed at the Interaction Region (IR) of the DAΦNE collider, is presented in this paper.

2. The SIDDHARTA-2 apparatus

The SIDDHARTA-2 apparatus is presently installed at the Interaction Region (IR) of the DAΦNE collider, in Frascati. The target consists of a cylindrical cell (144 mm in diameter and 125 mm in height) with walls made of a two-Kapton layer (150 μm thick) and aluminium supports. The target gas is fluxed inside the cylindrical cell. Outside, 384 Silicon Drift Detectors (SDDs) laterally surround the cylindrical target, for x-ray detection. SDDs provide a resolution of $157.8 \pm 0.3(stat)_{-0.2}^{+0.2}(syst)$ eV at 6.4 keV ($K^-^4\text{He}$, $3d \rightarrow 2p$ transition energy [5, 6]). Outside the cylindrical target and surrounding the SDDs, plastic scintillators read by pairs of Silicon Photo-Multipliers (SiPMs) are placed and used as a veto system for external background identification (VETO-2). The target cell, the SDDs and the VETO-2 are placed inside a vacuum chamber, which is kept at a pressure below 10^{-5} mbar. SDDs are cooled at about 120 K by a closed-cycle helium refrigerator system. Radially, outside the vacuum chamber, a second veto system consisting of 12 plastic scintillators read by pairs of Photo-Multipliers (PMTs) is installed for further external background reduction. Two plastic scintillators read by PMTs are placed one below and one above the DAΦNE Interaction Point (IP) and used to discriminate the back-to-back kaons directed to the SIDDHARTA-2 target, based on the time-of-flight (Kaon trigger system). A luminometer, consisting of two plastic scintillators read by pairs of photomultipliers, is placed on the horizontal side of the IP. A schematic drawing of the SIDDHARTA-2 apparatus is shown in Fig. 1. More detailed information on the elements of the SIDDHARTA-2 apparatus can be found in [3, 7]. The SIDDHARTINO experiment, consisting in a reduced SIDDHARTA-2 apparatus with only 64 SDDs installed, performed a precise measurement of gaseous $K^-^4\text{He}$ transitions to the $2p$ level [8, 9], thus confirming the expected performances, the excellent background rejection capacity of the SIDDHARTA-2 apparatus, and the validity of the data selection and analysis procedure [10].




Figure 1: A schematic drawing of the SIDDHARTA-2 apparatus installed at the DAΦNE Interaction Region at the INFN National Laboratories of Frascati.

3. Conclusions and future perspectives

The SIDDHARTA-2 experiment is presently installed at the DAΦNE e^-e^+ collider (see Fig. 2), and is ready to perform the first measurement of kaonic deuterium $2p \rightarrow 1s$ x-ray transitions. The data-taking campaign will start in 2023. The preliminary measurements performed by SIDDHARTINO and SIDDHARTA-2 on kaonic ^4He proved the high efficiency of the apparatus and the validity of the data analysis procedure.

Acknowledgments

We thank C. Capocchia from LNF-INFN and H. Schneider, L. Stohwasser, and D. Pristauz-Tel-snigg from Stefan Meyer-Institut for their fundamental contribution in designing and building the SIDDHARTA-2 setup. We thank as well the DAΦNE staff for the excellent working conditions and permanent support. Part of this work was supported by the Austrian Science Fund (FWF): [P24756-N20 and P33037-N]; the EXOTICA project of the Ministero degli Affari Esteri e della Cooperazione Internazionale, PO21MO03; the Croatian Science Foundation under the project IP-2018-01-8570; the EU STRONG-2020 project (Grant Agreement No. 824093); the EU Horizon 2020 project under the MSCA (Grant Agreement 754496); the Japan Society for the Promotion of Science JSPS KAKENHI Grant No. JP18H05402; the Polish Ministry of Science and Higher Education grant No. 7150/E-338/M/2018 and the Polish National Agency for Academic Exchange (grant no PPN/BIT/2021/1/00037). The authors acknowledge support from the SciMat and qLife Priority Research Areas budget under the program Excellence Initiative—Research University at the Jagiellonian University.



Figure 2: Picture of the SIDDHARTA-2 experiment presently installed in the DAΦNE Φ factory, at the National Laboratories of Frascati (LNF-INFN), in Italy.

References

- [1] C. Curceanu et al., *The modern era of light kaonic atom experiments* *Rev. Mod. Phys.* **91**, (2019) 025006
- [2] M. Bazzi et al., *A new measurement of kaonic hydrogen X-rays*, *Phys. Lett. B* **704** (2011) 113
- [3] F. Napolitano et al., *Kaonic atoms at the DAΦNE collider with the SIDDHARTA-2 experiment*, *Physica Scripta* **97** (2022) 8
- [4] M. Zobov et al., *Test of “Crab-Waist” Collisions at the DAΦNE Φ Factory*, *Phys. Rev. Lett.* **104** (2010) 174801
- [5] M. Miliucci et al., *Large area Silicon Drift Detectors system for high precision timed X-ray spectroscopy*, *Meas. Sci. Technol.* **33** (2022) 095502
- [6] M. Miliucci et al., *Silicon drift detectors system for high-precision light kaonic atoms spectroscopy*, *Meas. Sci. Technol.* **32** (2021) 095501
- [7] M. Tuchler et al., *Main Features of the SIDDHARTA-2 Apparatus for Kaonic Deuterium X-Ray Measurements*, *EPJ Web of Conf.* **262** (2022) 01016
- [8] D. L. Sirghi et al., *A new kaonic helium measurement in gas by SIDDHARTINO at the DAΦNE collider**, *J. Phys. G: Nucl. Part. Phys.* **49** (2022) 055106
- [9] D. L. Sirghi et al., *New measurements of kaonic helium-4 L-series X-rays yields in gas with the SIDDHARTINO setup*, *Nucl. Phys. A* (2022) 122567

- [10] F. Sgaramella et al., *The SIDDHARTA-2 calibration method for high precision kaonic atoms x-ray spectroscopy measurements*, *Phys. Scr.* **97** (2022) 114002