

Search for a dark photon in the  $\pi^0 \rightarrow e^+e^-\gamma$  decay

WASA-at-COSY Collaboration

P. Adlarson<sup>a</sup>, W. Augustyniak<sup>b</sup>, W. Bardan<sup>c</sup>, M. Bashkanov<sup>d,e</sup>, F.S. Bergmann<sup>f</sup>, M. Berłowski<sup>g</sup>, H. Bhatt<sup>h</sup>, A. Bondar<sup>i</sup>, M. Büscher<sup>j,k</sup>, H. Calén<sup>a</sup>, I. Ciepał<sup>c</sup>, H. Clement<sup>d,e</sup>, D. Coderre<sup>j,k,l</sup>, E. Czerwiński<sup>c</sup>, K. Demmich<sup>f</sup>, E. Doroshkevich<sup>d,e</sup>, R. Engels<sup>j,k</sup>, W. Erven<sup>m,k</sup>, W. Eyrych<sup>n</sup>, P. Fedorets<sup>j,k,o</sup>, K. Föhl<sup>p</sup>, K. Fransson<sup>a</sup>, F. Goldenbaum<sup>j,k</sup>, P. Goslawski<sup>f</sup>, A. Goswami<sup>q</sup>, K. Grigoryev<sup>j,k,r</sup>, C.-O. Gullström<sup>a</sup>, F. Hauenstein<sup>n</sup>, L. Heijkenkjöld<sup>a</sup>, V. Hejny<sup>j,k</sup>, F. Hinterberger<sup>s</sup>, M. Hodana<sup>c,j,k</sup>, B. Höistad<sup>a</sup>, A. Jany<sup>c</sup>, B.R. Jany<sup>c</sup>, L. Jarczyk<sup>c</sup>, T. Johansson<sup>a</sup>, B. Kamys<sup>c</sup>, G. Kemmerling<sup>m,k</sup>, F.A. Khan<sup>j,k</sup>, A. Khoukaz<sup>f</sup>, S. Kistryn<sup>c</sup>, J. Klaja<sup>c</sup>, H. Kleines<sup>m,k</sup>, D.A. Kirillov<sup>t</sup>, B. Kłos<sup>u</sup>, M. Krapp<sup>n</sup>, W. Krzemień<sup>c</sup>, P. Kulesa<sup>v</sup>, A. Kupść<sup>a,g,\*</sup>, A. Kuzmin<sup>i</sup>, K. Lalwani<sup>h,1</sup>, D. Lersch<sup>j,k</sup>, L. Li<sup>n</sup>, B. Lorentz<sup>j,k</sup>, A. Magiera<sup>c</sup>, R. Maier<sup>j,k</sup>, P. Marciniowski<sup>a</sup>, B. Mariański<sup>b</sup>, U.-G. Meißner<sup>j,k,w,s,x</sup>, M. Mikirtychiants<sup>j,k,l,r</sup>, H.-P. Morsch<sup>b</sup>, P. Moskal<sup>c</sup>, B.K. Nandi<sup>h</sup>, H. Ohm<sup>j,k</sup>, I. Ozerianska<sup>c</sup>, E. Perez del Rio<sup>d,e</sup>, N.M. Piskunov<sup>t</sup>, P. Pluciński<sup>a,2</sup>, P. Podkopał<sup>c,j,k</sup>, D. Prasuhn<sup>j,k</sup>, A. Pricking<sup>d,e</sup>, D. Pszczel<sup>a,g</sup>, K. Pysz<sup>v</sup>, A. Pyszniak<sup>a,c</sup>, C.F. Redmer<sup>a,3</sup>, J. Ritman<sup>j,k,l</sup>, A. Roy<sup>q</sup>, Z. Rudy<sup>c</sup>, S. Sawant<sup>h</sup>, S. Schadmand<sup>j,k</sup>, A. Schmidt<sup>n</sup>, T. Sefzick<sup>j,k</sup>, V. Serdyuk<sup>j,k,y</sup>, N. Shah<sup>h,4</sup>, B. Shwartz<sup>i</sup>, M. Siemaszko<sup>u</sup>, R. Siudak<sup>v</sup>, T. Skorodko<sup>d,e</sup>, M. Skurzok<sup>c</sup>, J. Smyrski<sup>c</sup>, V. Sopov<sup>o</sup>, R. Stassen<sup>j,k</sup>, J. Stepaniak<sup>g</sup>, E. Stephan<sup>u</sup>, G. Sterzenbach<sup>j,k</sup>, H. Stockhorst<sup>j,k</sup>, H. Ströher<sup>j,k</sup>, A. Szczurek<sup>v</sup>, T. Tolba<sup>j,k,5</sup>, A. Trzcíński<sup>b</sup>, R. Varma<sup>h</sup>, G.J. Wagner<sup>d,e</sup>, W. Węglorz<sup>u</sup>, A. Wirzba<sup>j,k,w</sup>, M. Wolke<sup>a</sup>, A. Wrońska<sup>c</sup>, P. Wüstner<sup>m,k</sup>, P. Wurm<sup>j,k</sup>, A. Yamamoto<sup>z</sup>, J. Zabierowski<sup>aa</sup>, M.J. Zieliński<sup>c</sup>, W. Zipper<sup>u</sup>, J. Złomańczuk<sup>a</sup>, P. Żuprański<sup>b</sup>, M. Żurek<sup>c</sup>

<sup>a</sup> Division of Nuclear Physics, Department of Physics and Astronomy, Uppsala University, Box 516, 75120 Uppsala, Sweden<sup>b</sup> Department of Nuclear Physics, National Centre for Nuclear Research, ul. Hoza 69, 00-681, Warsaw, Poland<sup>c</sup> Institute of Physics, Jagiellonian University, ul. Reymonta 4, 30-059 Kraków, Poland<sup>d</sup> Physikalisches Institut, Eberhard Karls Universität Tübingen, Auf der Morgenstelle 14, 72076 Tübingen, Germany<sup>e</sup> Kepler Center for Astro and Particle Physics, Eberhard Karls University Tübingen, Auf der Morgenstelle 14, 72076 Tübingen, Germany<sup>f</sup> Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, Wilhelm-Klemm-Str. 9, 48149 Münster, Germany<sup>g</sup> High Energy Physics Department, National Centre for Nuclear Research, ul. Hoza 69, 00-681, Warsaw, Poland<sup>h</sup> Department of Physics, Indian Institute of Technology Bombay, Powai, Mumbai 400076, Maharashtra, India<sup>i</sup> Budker Institute of Nuclear Physics of SB RAS, Academician Lavrentyev 11, Novosibirsk, 630090, Russia<sup>j</sup> Institut für Kernphysik, Forschungszentrum Jülich, 52425 Jülich, Germany<sup>k</sup> Jülich Center for Hadron Physics, Forschungszentrum Jülich, 52425 Jülich, Germany<sup>l</sup> Institut für Experimentalphysik I, Ruhr-Universität Bochum, Universitätsstr. 150, 44780 Bochum, Germany<sup>m</sup> Zentralinstitut für Engineering, Elektronik und Analytik, Forschungszentrum Jülich, 52425 Jülich, Germany<sup>n</sup> Physikalisches Institut, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erwin-Rommel-Str. 1, 91058 Erlangen, Germany<sup>o</sup> Institute for Theoretical and Experimental Physics, State Scientific Center of the Russian Federation, Bolshaya Chermushkinskaya 25, 117218 Moscow, Russia<sup>p</sup> II. Physikalisches Institut, Justus-Liebig-Universität Gießen, Heinrich-Buff-Ring 16, 35392 Giessen, Germany<sup>q</sup> Department of Physics, Indian Institute of Technology Indore, Khandwa Road, Indore 452017, Madhya Pradesh, India<sup>r</sup> High Energy Physics Division, Petersburg Nuclear Physics Institute, Orlova Rosh 2, Gatchina, Leningrad district 188300, Russia<sup>s</sup> Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität Bonn, Nuffallee 14-16, 53115 Bonn, Germany<sup>t</sup> Veksler and Baldin Laboratory of High Energy Physics, Joint Institute for Nuclear Physics, Joliot-Curie 6, 141980 Dubna, Moscow region, Russia<sup>u</sup> August Chelkowski Institute of Physics, University of Silesia, Uniwersytecka 4, 40-007, Katowice, Poland<sup>v</sup> The Henryk Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences, 152 Radzikowskiego St., 31-342 Kraków, Poland<sup>w</sup> Institute for Advanced Simulation, Forschungszentrum Jülich, 52425 Jülich, Germany<sup>x</sup> Bethe Center for Theoretical Physics, Rheinische Friedrich-Wilhelms-Universität Bonn, 53115 Bonn, Germany<sup>y</sup> Dzhelapov Laboratory of Nuclear Problems, Joint Institute for Nuclear Physics, Joliot-Curie 6, 141980 Dubna, Moscow region, Russia<sup>z</sup> High Energy Accelerator Research Organisation KEK, Tsukuba, Ibaraki 305-0801, Japan<sup>aa</sup> Department of Cosmic Ray Physics, National Centre for Nuclear Research, ul. Uniwersytecka 5, 90-950 Łódź, Poland

## ARTICLE INFO

## Article history:

Received 3 April 2013

Received in revised form 7 July 2013

Accepted 21 August 2013

Available online 28 August 2013

Editor: V. Metag

## Keywords:

Dark forces

Gauge vector boson

## ABSTRACT

The presently world largest data sample for  $\pi^0 \rightarrow \gamma e^+ e^-$  decays studies containing nearly  $5 \times 10^5$  events was collected using the WASA detector at COSY. A search for a dark photon  $U$  produced in the  $\pi^0 \rightarrow \gamma U \rightarrow \gamma e^+ e^-$  decay from the  $pp \rightarrow pp\pi^0$  reaction was carried out. An upper limit on the square of the  $U - \gamma$  mixing strength parameter  $\epsilon^2$  of  $5 \times 10^{-6}$  at 90% CL was obtained for the mass range  $20 \text{ MeV} < M_U < 100 \text{ MeV}$ . This result together with other recent experimental limits significantly reduces the  $M_U$  vs.  $\epsilon^2$  parameter space which could explain the presently seen deviation between the Standard Model prediction and the direct measurement of the anomalous magnetic moment of the muon.

© 2013 Elsevier B.V. All rights reserved.

## 1. Introduction

Decays of neutral pseudoscalar mesons into a lepton–antilepton pair and a photon,  $P \rightarrow l^+ l^- \gamma$ , are among the processes to search for a new light vector boson connected with dark gauge forces [1–3]. An extra  $U(1)$  boson is postulated in most extensions of the Standard Model. Recent interest in searches of a light vector boson, in the  $\mathcal{O}(\text{MeV} - \text{GeV})$  mass range, is motivated by astrophysics observations such as the positron and/or electron excesses observed by PAMELA [4], ATIC [5] and H.E.S.S. [6] as well as the narrow 0.511 MeV  $\gamma$  ray emission from the galactic bulge observed by INTEGRAL [7].

In one of the simplest scenarios dark matter particles belonging to an additional abelian gauge symmetry are added to the Standard Model (SM). The new symmetry leaves the SM particles unchanged [8,9,3,10]. The associated gauge boson can communicate with the SM through a small mixing in the kinetic term of the QED Lagrangian [11]:

$$\mathcal{L}_{\text{mix}} = -\frac{\epsilon}{2} F_{\mu\nu}^{\text{QED}} F_{\text{dark}}^{\mu\nu} \quad (1)$$

where  $\epsilon$  is the mixing parameter. The gauge boson  $U$  (also  $A'$ ,  $\gamma'$  or  $Z'_d$ ) is often called a *dark photon* since it can mix with the photon in all processes (examples are shown in Figs. 1b and 1c). Phenomenological arguments [12–14] suggest that the  $\epsilon$  parameter must be of the order of  $10^{-4} - 10^{-2}$  and the boson mass  $M_U$  below 2 GeV. This estimate is also supported by the astrophysical observations and the constraints imposed by precision measurements such as the anomalous magnetic moments ( $g - 2$ ) of muon and electron [15]. The contribution of the  $U$  boson to the  $(g - 2)_l$  ( $l = e, \mu$ ) (Fig. 1c) is given in [15] by:

$$\Delta(g - 2)_l = \frac{\alpha \epsilon^2}{\pi} \int_0^1 dz \frac{2m_l^2 z(1-z)^2}{m_l^2(1-z)^2 + M_U^2 z} \quad (2)$$

Investigations of the  $M_U$  vs.  $\epsilon^2$  parameter space corresponding to the experimentally preferred  $(g - 2)_\mu$  value (shifted  $+3.6\sigma$  with respect to the SM value [16–18]) are therefore of great importance.

\* Corresponding author at: Division of Nuclear Physics, Department of Physics and Astronomy, Uppsala University, Box 516, 75120 Uppsala, Sweden.

E-mail address: andrzej.kupsc@physics.uu.se (A. Kupść).

<sup>1</sup> Present address: Department of Physics and Astrophysics, University of Delhi, Delhi 110007, India.

<sup>2</sup> Present address: Department of Physics, Stockholm University, Roslagstullsbacken 21, AlbaNova, 10691 Stockholm, Sweden.

<sup>3</sup> Present address: Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, Johann-Joachim-Becher Weg 45, 55128 Mainz, Germany.

<sup>4</sup> Present address: Department of Physics and Astronomy, University of California, Los Angeles, CA 90045, USA.

<sup>5</sup> Present address: Albert Einstein Center for Fundamental Physics, Fachbereich Physik und Astronomie, Universität Bern, Sidlerstr. 5, 3012 Bern, Switzerland.

For a  $U$  boson with mass less than twice the muon mass the total decay width is for all practical purposes (neglecting higher-order electric, tiny weak interaction contributions from the  $U$  boson –  $Z_0$  coupling, and the decay to light dark scalars and/or fermions) given by [19,20]:

$$\Gamma_U = \Gamma_{U \rightarrow e^+ e^-} = \frac{1}{3} \alpha \epsilon^2 M_U \sqrt{1 - \frac{4m_e^2}{M_U^2}} \left( 1 + \frac{2m_e^2}{M_U^2} \right), \quad (3)$$

where  $m_e$  is the electron mass.

Such a light  $U$  boson can be directly produced in particle accelerators, see e.g. Refs. [19–27]. The idea is to search for narrow structures in the invariant mass spectrum of the lepton–antilepton pair.

The  $M_U$  vs.  $\epsilon^2$  region corresponding to the measured  $(g - 2)_\mu$  value  $\pm 2\sigma$  is covered by the data from the BABAR [28], MAMI A1 [29], KLOE-2 [30] and APEX [31] experiments for  $M_U$  masses above 100 MeV. On the lower end this preferred region is excluded by the  $(g - 2)_e$  value for  $M_U < 30 \text{ MeV}$  [32,33]. In addition,  $\epsilon^2$  regions below  $10^{-12}$  are excluded by experiments which are sensitive to lepton pairs from displaced secondary vertices ( $\tau_U > 10^{-11} \text{ s}$ ) [34–36].


Our experiment aims at searching for a short-lived  $U$  boson in the  $\pi^0$  Dalitz decay,  $\pi^0 \rightarrow e^+ e^- \gamma$ , covering the range preferred by the experimental value of  $(g - 2)_\mu$  for  $20 \text{ MeV} < M_U < 100 \text{ MeV}$ . In this region, for  $\epsilon^2 > 10^{-6}$  the average distance passed by a boson emitted from a low energy  $\pi^0$  decay should be less than a millimeter. The best limit from a previous  $\pi^0 \rightarrow e^+ e^- \gamma$  experiment with the origin of the  $e^+ e^-$  pair close to the production vertex was obtained by the SINDRUM collaboration more than twenty years ago [37,38]. The SINDRUM result is based on a sample of 98 400  $\pi^0 \rightarrow e^+ e^- \gamma$  decays with  $e^+ e^-$  invariant masses above 25 MeV.

## 2. The experiment

The WASA detector setup was built and first used at CELSIUS in Uppsala and moved to COSY (COoler SYNchrotron) Jülich in the Summer of 2005 [39]. The detector was designed and optimized for studies of rare  $\pi^0$  meson decays produced in  $pp \rightarrow pp\pi^0$  reaction [40]. It consists of three main components:

The Forward Detector (FD) – covering scattering angles in the  $3^\circ - 18^\circ$  range used for tagging and triggering of meson production, the Central Detector (CD) – used for measuring meson decay products, and the pellet target system. The target beam consists of 20–30  $\mu\text{m}$  diameter pellets of hydrogen, providing an areal target density in the order of  $10^{15} \text{ atoms/cm}^2$ . The diameter of the pellet beam is  $\sim 3.8 \text{ mm}$ .

The CD surrounds the interaction region and is designed to detect and identify photons, electrons, and charged pions. It consists of an inner drift chamber (MDC), a superconducting solenoid providing the magnetic field for momentum determination, a barrel of



**Fig. 1.** Feynman diagrams for a) the lowest order electromagnetic  $\pi^0 \rightarrow e^+e^-\gamma$  decay and a possible contribution of  $U$  vector boson to: b)  $\pi^0 \rightarrow e^+e^-\gamma$  and c) lepton  $g-2$ .

thin plastic scintillators (PS) for particle identification and triggering, and an electromagnetic calorimeter. The amount of structural material is kept to a minimum to reduce the amount of secondary interactions outside of the detector sensitive volumes. The beryllium beam pipe (diameter 6 cm) wall is 1.2 mm thick and the material of the superconducting solenoid corresponds to 0.18 radiation lengths.


The FD allows identification and reconstruction of protons from the  $pp \rightarrow pp\pi^0$  reaction close to threshold. The track coordinates are provided by four sets of straw proportional chambers. Kinetic energies are reconstructed using the  $\Delta E$  information in layers of plastic scintillators of different thickness. In addition, the signals are used for triggering. The kinetic energy,  $T$ , of the protons can be reconstructed with a resolution of  $\sigma(T)/T \sim 1.5\text{--}3\%$  for kinetic energies below 400 MeV.

The results presented here are based on data collected during one-week WASA-at-COSY run carried out in 2010. The  $\pi^0$  mesons were produced in proton–proton interactions at a kinetic beam energy of 550 MeV. The beam energy corresponds to the center-of-mass excess energy of 122 MeV with respect to  $pp\pi^0$  threshold (i.e. below two pion production thresholds) with a cross section of 1.12 mb [41]. The maximum scattering angle of the outgoing protons for the reaction is  $45^\circ$ . For detection and for triggering purposes the phase space of the  $pp \rightarrow pp\pi^0$  reaction can be divided into three regions:

- (1) Both protons are measured in the FD. This corresponds to a geometrical acceptance of 19%.
- (2) One proton is measured in the FD and one in the forward part of the PS (scattering angles  $20^\circ\text{--}40^\circ$ ). This corresponds to a geometrical acceptance of 42%.
- (3) Both protons are registered in the PS. This corresponds to a geometrical acceptance of 21%.


Case (1) allows the definition of the most selective trigger condition and the best resolution in the missing mass with respect to the two protons. Therefore, the main trigger for the experiment required two tracks in the FD. The protons from the  $pp \rightarrow pp\pi^0$  reaction have a maximum kinetic energy of 350 MeV and are mostly stopped in the FD. This allows the inclusion of a veto from a thin plastic detector layer placed at the far end of the FD into the trigger condition. In addition, two hits in the central part of the PS (scattering angles  $45^\circ\text{--}135^\circ$ ) were required, aiming to select the electron–positron pair. An additional, scaled down, trigger based on case (2) was used in parallel. The WASA-at-COSY data acquisition system allowed the collection of more than  $10^4$  events per second and the luminosity was set to optimize the conditions for the main trigger. The integrated luminosity of the run was about  $0.55 \text{ pb}^{-1}$ .

The data quality is illustrated by analysis of the main trigger data sample and requesting in the analysis two identified (using  $\Delta E/\Delta E$  method) FD proton tracks. An electron–positron pair is selected by requiring two oppositely curved tracks in the MDC with scattering angles between  $40^\circ$  and  $140^\circ$ . A photon hit cluster in the calorimeter with an energy deposit above 20 MeV is also requested. The missing mass squared with respect to two protons



**Fig. 2.** Detector performance plots for a data sample with two reconstructed protons, an  $e^+e^-$  pair and a photon. a) Distribution of the missing mass squared with respect to the two protons registered in the FD before electron identification. Experimental data (black points); simulations:  $\pi^0 \rightarrow e^+e^-\gamma$  and  $\pi^0 \rightarrow \gamma\gamma$  (broken line), random coincidences of two events (dotted line), and the sum (solid line). b) Distribution of  $MM(pp)$  after electron identification; experimental data (black points) and sum of Monte Carlo simulations (solid line). c) The reconstructed invariant mass of the  $e^+e^-\gamma$  system after particle identification cut.

( $MM^2(pp)$ ) for the above selection is shown in Fig. 2a. In addition to the  $pp \rightarrow pp\pi^0$  reaction signal one sees also a contribution due to random coincidences of  $pp \rightarrow pp$  and  $pp \rightarrow pn\pi^+$  reactions. This background is effectively suppressed by including electron and positron identification using the reconstructed momentum and the energy deposit in the calorimeter. The corresponding  $MM(pp)$  plot after this cut is shown in Fig. 2b. The  $\pi^0 \rightarrow e^+e^-\gamma$  decay is independently identified from the invariant mass of the decay products  $IM(e^+e^-\gamma)$  (calculated assuming the tracks originate at the beam target crossing) shown in Fig. 2c. The data are well described by



**Fig. 3.** The reconstructed  $e^+e^-$  invariant mass  $q = IM(e^+e^-)$ : a) before and b) after the cuts for reducing the conversion background. The experimental data are denoted by black points. Results of simulations for  $\pi^0 \rightarrow \gamma\gamma$  (blue line) and  $\pi^0 \rightarrow e^+e^-\gamma$  (green line) decays are normalized according to the known branching ratios. The normalization of random coincidences (dotted line) was fitted in order to reproduce the  $IM(e^+e^-) > 150$  MeV range. The sum of all simulated contributions is given by the red line. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this Letter.)

a simulation of  $pp \rightarrow pp\pi^0$  with  $\pi^0 \rightarrow e^+e^-\gamma$  and  $\pi^0 \rightarrow \gamma\gamma$  decays, where in the latter case one of the two photons converts in the beryllium beam tube.

For Monte Carlo simulations, angular distributions for the  $pp \rightarrow pp\pi^0$  reaction from [41] were used in the event generation. The  $\pi^0 \rightarrow e^+e^-\gamma$  decay is generated using the lowest order QED matrix element squared:

$$|A|^2 = \Gamma_{\gamma\gamma} 16\pi^3 M \frac{\alpha}{\pi} \frac{1}{q^2} \left(1 - \frac{q^2}{M^2}\right)^2 \times \left(1 + \cos^2 \theta^* + \frac{4m_e^2}{q^2} \sin^2 \theta^*\right) |F(q^2)|^2 \quad (4)$$


where  $\theta^*$  is the angle of  $e^+$  in the dilepton rest frame with respect to the dilepton momentum in the overall  $\pi^0$  decay system,  $M$  and  $m_e$  are  $\pi^0$  and  $e^\pm$  masses respectively,  $\Gamma_{\gamma\gamma}$  is the partial  $\pi^0 \rightarrow \gamma\gamma$  decay width, and  $F(q^2)$  (with  $q^2$  the squared momentum transfer of the off-shell photon) is the  $\pi^0$  transition form factor. The form factor close to  $q^2 = 0$  is parametrized as:  $F(q^2) = 1 + aq^2/M^2$ . The value of the dimensionless linear coefficient  $a$  is  $0.032 \pm 0.004$  [42].

The matrix element from Eq. (4) leads to the following unperturbed  $d\Gamma/dq$  distribution [43] for the standard lowest order electromagnetic decay  $\pi^0 \rightarrow e^+e^-\gamma$  of Fig. 1a:

$$\frac{d\Gamma}{dq} = \Gamma_{\gamma\gamma} \frac{4\alpha}{3\pi} \frac{1}{q} \sqrt{1 - \frac{4m_e^2}{q^2}} \left(1 + \frac{2m_e^2}{q^2}\right) \times \left(1 - \frac{q^2}{M^2}\right)^3 |F(q^2)|^2. \quad (5)$$

### 3. Data analysis


The first stage of data analysis is to extract a clean signal of  $\pi^0 \rightarrow e^+e^-\gamma$  decays. The results shown in the previous section suggest that in  $pp$  interactions at 550 MeV electron-positron pairs




**Fig. 4.** Distribution of the distance  $R$  between the COSY beam axis and the reconstructed point of closest approach of  $e^+$  and  $e^-$  tracks: experimental data (black crosses); simulations for  $\pi^0 \rightarrow \gamma\gamma$  (blue line), the  $\pi^0 \rightarrow e^+e^-\gamma$  decay (green line), and the sum of the two contributions (red line). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this Letter.)

come nearly exclusively from the  $\pi^0$  meson decays. Therefore, in order to maximize the yield of the  $\pi^0 \rightarrow \gamma e^+e^-$  events we use an inclusive data sample requesting events with (i) at least one proton identified in the FD, (ii) an  $e^+e^-$  pair identified in the CD. There is no request of an additional photon cluster and we have included events from both triggers corresponding to phase space regions (1) and (2). The distribution of the reconstructed invariant mass of the electron-positron pair,  $q = IM(e^+e^-)$ , is shown in Fig. 3a. This spectrum is well described by the sum of  $\pi^0 \rightarrow e^+e^-\gamma$  and  $\pi^0 \rightarrow \gamma\gamma$  (with photon conversion). The data sample contains  $1.8 \times 10^6$  reconstructed events.

The  $\pi^0 \rightarrow \gamma\gamma$  events are efficiently removed by a condition on the reconstructed position of the  $e^+e^-$  vertex. Fig. 4 shows the distance ( $R$ ) of the reconstructed vertex from the COSY beam axis. The contributions of the  $\pi^0 \rightarrow \gamma\gamma$  and  $\pi^0 \rightarrow e^+e^-\gamma$  decays, simulated according to the known branching ratios, are in



**Fig. 6.** Difference between the reconstructed  $e^+e^-$  invariant mass distribution and the sum of all simulated contributions (black points). The resolution and sensitivity for a hypothetical decay  $U \rightarrow e^+e^-$  are illustrated by the superimposed red histograms. They represent the signals expected for the  $\pi^0 \rightarrow U\gamma \rightarrow e^+e^-\gamma$  process with  $U$  boson masses of  $M_U = 30, 50, 70$  and  $90$  MeV and  $BR(\pi^0 \rightarrow U\gamma) = 10^{-4}$  (the corresponding  $\epsilon$  values are: 0.0077, 0.0088, 0.0113, and 0.0169 respectively).



**Fig. 5.** Correlation between  $R$  and  $IM_b$  variables for the experimental data. The selection cut is shown by the diagonal line. The events below the line mainly come from photon conversions in the beam pipe.

very good agreement with the observed distribution and they are well separated. In order to further reduce the external conversion background one uses the invariant mass of the  $e^+e^-$  calculated from the momentum directions at the points where the tracks intersect the beam tube,  $IM_b$ , shown in Fig. 5. The selection cut is performed in the  $IM_b$  vs.  $R$  plane (Fig. 5). The cut removes 98% of the  $\pi^0 \rightarrow \gamma\gamma$  events which contribute to  $IM(e^+e^-)$  distribution due to conversion.

The finally reconstructed  $dN/dq$  distribution, containing nearly  $5 \times 10^5$  entries, is shown in Fig. 3b. It is well described by the simulations of the  $\pi^0 \rightarrow e^+e^-\gamma$  decay channel alone with a very small (approx. 3000 events) admixture of background from the  $\pi^0 \rightarrow \gamma\gamma$  decay. The data in this work represent the world largest data sample of  $\pi^0 \rightarrow e^+e^-\gamma$  events, which is almost an order of magnitude larger than the sample used for the previously published results from the SINDRUM experiment [37,44].

### 3.1. Upper limit for the $BR(\pi^0 \rightarrow \gamma(U \rightarrow e^+e^-))$

A distinctive feature of the expected signal of the decay  $\pi^0 \rightarrow \gamma(U \rightarrow e^+e^-)$  (Fig. 1b) is the appearance of a narrow peak (the width being given by the detector resolution) in the invariant mass distribution of the electron-positron pair at the  $U$  boson mass. The electrodynamic process  $\pi^0 \rightarrow \gamma^*\gamma \rightarrow e^+e^-\gamma$  (Fig. 1a) both represents the irreducible background and is used for normalization. Due to the expected small decay width of the  $U$  boson the interference term is negligible and the signal from the  $U$  boson can be tested by constructing an incoherent sum of the two contributions.

The experimental data are described well by the simulation based on Eq. (4) alone as shown in Fig. 3. The difference between reconstructed experimental  $q$  distribution and the sum of all sim-

ulated contributions is given in Fig. 6. The errors include both statistical uncertainties of the data sample as well as the systematical ones due to the simulation of the detector response. In addition there are superimposed five example distributions corresponding to the  $\pi^0 \rightarrow U\gamma \rightarrow e^+e^-\gamma$  process for  $U$  boson masses of 30, 50, 70 and 90 MeV respectively, assuming  $BR(\pi^0 \rightarrow U\gamma) = 10^{-4}$ . The plots illustrate both the resolution and the efficiency expected for the signal. The structure at 60 MeV is most likely due to a small residual of the conversion events which are not yet understood by MC.

For a given value of the  $U$  boson mass corresponding to the range of the  $k$ th bin of the invariant mass spectrum ( $q_k < M_U < q_k + \Delta q$ , with  $\Delta q = 1$  MeV the width of the histogram bin) the number of events in the  $i$ th bin of the reconstructed electron-positron invariant mass distribution,  $N_i$ , can be described in the following form:

$$N_i/N_{Tot} = \frac{1}{\Gamma} \sum_j S_{ij}\eta_j\nu_j + S_{ik}\eta_k\beta. \quad (6)$$

The first term in Eq. (6) represents the contribution from the Dalitz decay and the second term from the hypothetical  $\pi^0 \rightarrow \gamma(U \rightarrow e^+e^-)$  decay chain. Indices  $j$  and  $k$  label the *true, unperturbed* distributions and  $i$  the reconstructed  $q$  histogram.  $N_{Tot}$  is total number of produced  $\pi^0$  mesons,  $1/\Gamma$  is the  $\pi^0$  life time and  $\eta_j$  is the efficiency.  $S_{ij}$  is the normalized smearing matrix (for each  $j$ :  $\sum_i S_{ij} = 1$ ),  $\nu_j$  is the unperturbed  $d\Gamma/dq$  distribution for the  $\pi^0 \rightarrow e^+e^-\gamma$  decay (Eq. (5) and Fig. 1a) integrated over bin  $j$ :

$$\nu_j \equiv \int_{q_j}^{q_j+\Delta q} \frac{d\Gamma}{dq} dq, \quad (7)$$

and  $\beta$  is  $BR(\pi^0 \rightarrow \gamma(U \rightarrow e^+e^-))$ . The efficiency and the smearing matrix was obtained from the detector simulation. The  $U$  boson decay mechanism in diagram Fig. 1b implies that the efficiencies as a function of  $\cos\theta^*$  are identical to the ones of the  $\pi^0 \rightarrow e^+e^-\gamma$  decay with  $q = M_U$ . Note that for the quoted values of the branching ratios the intrinsic width (3) of the  $U$  boson would be in the eV range and thus very much smaller than the experimental bin size.

The upper limits for the  $U$  boson branching ratios,  $\beta$ , as a function of  $M_U$  were obtained by repeating for all bins (index  $k$  in Eq. (6)), corresponding to the  $20 \text{ MeV} < M_U < 100 \text{ MeV}$  range, the least square fits of Eq. (6) to the experimental  $q$  distribution. The results of the unconstrained fits yield estimators of  $\beta$  values and their standard deviations, which have to a good accuracy asymptotic Gaussian distributions. Finally we construct the upper limits using prescriptions from Ref. [45] taking into account the fact that




Fig. 7. A 90% C.L. upper limit (smoothed) for the  $BR(\pi^0 \rightarrow \gamma U)$  from this Letter (solid line) compared to the result of the SINDRUM experiment [37] (dotted line).




Fig. 8. Summary of the 90% CL upper limits for the mixing parameter  $\epsilon^2$  from WASA-at-COSY (red solid line) compared to SINDRUM  $\pi^0 \rightarrow e^+e^-\gamma$  [37] (dotted line) and recent combined KLOE  $\phi \rightarrow \eta e^+e^-$  [47] (dashed dotted) upper limits. The long respectively short dashed lines (and the corresponding hatched areas) are the upper limits derived from the muon and the electron  $g-2$  [32]. In addition the gray area represents the  $\pm 2\sigma$  preferred band around the present value of the muon  $g-2$ .

$\beta$  is a non-negative parameter since the  $U$  boson contribution is added incoherently here. Fig. 7 shows the 90% C.L. upper limits for the branching ratio of  $\pi^0 \rightarrow \gamma(U \rightarrow e^+e^-)$  decay as a function of the assumed value of  $M_U$ . This result is compared to that obtained from the SINDRUM data [37].

The branching ratio of  $\pi^0 \rightarrow \gamma U$  is related to  $\epsilon^2$  by [46,21]:

$$\frac{\Gamma(\pi^0 \rightarrow \gamma U)}{\Gamma(\pi^0 \rightarrow \gamma \gamma)} = 2\epsilon^2 |F(M_U^2)|^2 \left(1 - \frac{M_U^2}{M^2}\right)^3. \quad (8)$$

The resulting upper limits for the  $\epsilon^2$  parameter is shown in Fig. 8 and compared with other experiments.

The recent limits for the electron  $g-2$  are taken from recent QED calculations Refs. [33,32] and a measurement of alpha in atomic physics [48]. Our upper limit improves the recent combined KLOE limits [47] at low  $M_U$ . We use a disparate experimental setup and different meson decay as source of  $e^+e^-$  pairs. Together the data significantly reduce the parameter space for mass and mixing strength of a hypothetical dark photon  $U$ , if the latter is assumed to account for the presently seen deviation between the Standard Model prediction and the experimental value of the muon anomalous magnetic moment. The experiment presented in the Letter if repeated with an order of magnitude larger statistics would cover the remaining part of this region of interest. The collected data can also be used to determine the  $\pi^0$  transition form factor.

## Acknowledgements

This work was supported in part by the EU Integrated Infrastructure Initiative HadronPhysics Project under contract number RII3-CT-2004-506078; by the European Commission under the 7th Framework Programme through the ‘Research Infrastructures’ action of the ‘Capacities’ Programme, Call: FP7-INFRASTRUCTURES-2008-1, Grant Agreement N. 227431; by the Polish National Science Centre through the Grants Nos. 86/2/N-DFG/07/2011/0, 0320/B/H03/2011/40, 2011/01/B/ST2/00431, 2011/03/B/ST2/01847, 0312/B/H03/2011/40 and Foundation for Polish Science. We gratefully acknowledge the support given by the Swedish Research Council, the Knut and Alice Wallenberg Foundation, and the Forschungszentrum Jülich FFE Funding Program of the Jülich Center for Hadron Physics.

We would like to thank Simona Giovannella for providing the KLOE data points. The authors thank technical staff at Forschungszentrum Jülich for support in preparation of and during the experiment.

This work is part of the PhD thesis of C.-O. Gullström.

## References

- [1] P. Fayet, Phys. Lett. B 95 (1980) 285.
- [2] M. Dobroliubov, A.Y. Ignatiev, Phys. Lett. B 206 (1988) 346.
- [3] C. Boehm, P. Fayet, Nucl. Phys. B 683 (2004) 219, arXiv:hep-ph/0305261.
- [4] O. Adriani, et al., PAMELA Collaboration, Nature 458 (2009) 607, arXiv:0810.4995.
- [5] J. Chang, et al., Nature 456 (2008) 362.
- [6] F. Aharonian, et al., H.E.S.S. Collaboration, Phys. Rev. Lett. 101 (2008) 261104, arXiv:0811.3894.
- [7] P. Jean, et al., Astron. Astrophys. 407 (2003) L55, arXiv:astro-ph/0309484.
- [8] P. Fayet, Nucl. Phys. B 187 (1981) 184.
- [9] B. Holdom, Phys. Lett. B 166 (1986) 196.
- [10] M. Pospelov, A. Ritz, M.B. Voloshin, Phys. Lett. B 662 (2008) 53, arXiv:0711.4866.
- [11] L. Okun, Sov. Phys. JETP 56 (1982) 502.
- [12] N. Arkani-Hamed, et al., Phys. Rev. D 79 (2009) 015014, arXiv:0810.0713.
- [13] K.R. Dienes, C.F. Kolda, J. March-Russell, Nucl. Phys. B 492 (1997) 104, arXiv:hep-ph/9610479.
- [14] M. Goodsell, et al., J. High Energy Phys. 0911 (2009) 027, arXiv:0909.0515.
- [15] M. Pospelov, Phys. Rev. D 80 (2009) 095002, arXiv:0811.1030.
- [16] G. Bennett, et al., Muon G-2 Collaboration, Phys. Rev. D 73 (2006) 072003, arXiv:hep-ex/0602035.
- [17] M. Davier, et al., Eur. Phys. J. C 71 (2011) 1515, arXiv:1010.4180.
- [18] K. Hagiwara, et al., J. Phys. G 38 (2011) 085003, arXiv:1105.3149.
- [19] J.D. Bjorken, et al., Phys. Rev. D 80 (2009) 075018, arXiv:0906.0580.
- [20] B. Batell, M. Pospelov, A. Ritz, Phys. Rev. D 79 (2009) 115008, arXiv:0903.0363.
- [21] M. Reece, L.T. Wang, J. High Energy Phys. 0907 (2009) 051, arXiv:0904.1743.
- [22] M. Freytsis, G. Ovanesyan, J. Thaler, J. High Energy Phys. 1001 (2010) 111, arXiv:0909.2862.
- [23] N. Borodatchenkova, D. Choudhury, M. Drees, Phys. Rev. Lett. 96 (2006) 141802, arXiv:hep-ph/0510147.
- [24] P.f. Yin, J. Liu, S.h. Zhu, Phys. Lett. B 679 (2009) 362, arXiv:0904.4644.
- [25] R. Essig, P. Schuster, N. Toro, Phys. Rev. D 80 (2009) 015003, arXiv:0903.3941.
- [26] M. Baumgart, et al., J. High Energy Phys. 0904 (2009) 014, arXiv:0901.0283.
- [27] H.B. Li, T. Luo, Phys. Lett. B 686 (2010) 249, arXiv:0911.2067.
- [28] B. Aubert, et al., BABAR Collaboration, Phys. Rev. Lett. 103 (2009) 081803, arXiv:0905.4539.
- [29] H. Merkel, et al., A1 Collaboration, Phys. Rev. Lett. 106 (2011) 251802, arXiv:1101.4091.
- [30] F. Archilli, et al., Phys. Lett. B 706 (2012) 251, arXiv:1110.0411.
- [31] S. Abrahamyan, et al., APEX Collaboration, Phys. Rev. Lett. 107 (2011) 191804, arXiv:1108.2750.
- [32] M. Endo, K. Hamaguchi, G. Mishima, Phys. Rev. D 86 (2012) 095029, arXiv:1209.2558.
- [33] T. Aoyama, et al., Phys. Rev. Lett. 109 (2012) 111807, arXiv:1205.5368.
- [34] C. Amsler, et al., Crystal Barrel Collaboration, Phys. Lett. B 333 (1994) 271.
- [35] J. Altegoer, et al., NOMAD Collaboration, Phys. Lett. B 428 (1998) 197, arXiv:hep-ex/9804003.
- [36] S. Gninenko, Phys. Rev. D 85 (2012) 055027, arXiv:1112.5438.
- [37] R. Meijer Drees, et al., SINDRUM I Collaboration, Phys. Rev. Lett. 68 (1992) 3845.
- [38] S. Gninenko, arXiv:1301.7555, 2013.

- [39] H.H. Adam, et al., WASA-at-COSY Collaboration, arXiv:nucl-ex/0411038, 2004.
- [40] C. Bargholtz, et al., CELSIUS/WASA Collaboration, *Nucl. Instrum. Methods A* 594 (2008) 339, arXiv:0803.2657.
- [41] G. Rappenecker, et al., *Nucl. Phys. A* 590 (1995) 763.
- [42] J. Beringer, et al., Particle Data Group, *Phys. Rev. D* 86 (2012) 010001.
- [43] L. Landsberg, *Phys. Rep.* 128 (1985) 301.
- [44] R. Meijer Drees, et al., SINDRUM I Collaboration, *Phys. Rev. D* 45 (1992) 1439.
- [45] G. Cowan, et al., *Eur. Phys. J. C* 71 (2011) 1554, arXiv:1007.1727.
- [46] B. Batell, M. Pospelov, A. Ritz, *Phys. Rev. D* 80 (2009) 095024, arXiv:0906.5614.
- [47] D. Babusci, et al., KLOE-2 Collaboration, *Phys. Lett. B* 720 (2013) 111, arXiv:1210.3927.
- [48] R. Bouchendira, et al., *Phys. Rev. Lett.* 106 (2011) 080801.