




Positronium atom : a purely leptonic object to perform the tests on discrete symmetries

Sushil K. Sharma on behalf of the J-PET collaboration




2012



Characterize scintillator properties
Energy, time resolution, hit time,...


2014



24 strips

Data acquisition for multi-modules

192 strips



2016

Current version

Tests on discrete symmetries



24 modules, each with 13 scintillators

2018



Modular PET – ready for first data campaign

2-D front view J-PET



192 plastic Scintillators

Arranged axially in **3 layers**

Key features :

- ✓ Trigger less and reconfigurable **DAQ**
- ✓ **TOT** as measure of energy deposition


- ✓ *Good angular resolution and small light attenuation*
- ✓ *Superior time properties and lower pile-ups*



TOT as a response of energy deposition by incident photon



Sushil K. Sharma




Signals from each photomultiplier are probed at four thresholds.

$$TOT = TOT_{\text{sum}A} + TOT_{\text{sum}B}$$





Recipe to establish relationship between TOT and Energy dep(scatt. Ang.)



What we have :

- ✓ Measured **TOT** values
- ✓ Hit positions and time of primary and scattered photon gives access to the θ values

What is required :

- ✦ **Identify** the origin of **incident photon**

$$E_{\text{dep}} = f(E_{\text{inc}}, \theta)$$






Analyzed : 3 Hit Events


- In order to study *scattering of a photon*, two hits are sufficient.
- 3rd Hit allows to use additional constraints to conjecture and tag the photon of different energy/origin

$$\text{Scatter test} = \text{time}_{\text{measured}} - \text{time}_{\text{calculated}}$$



Case 1

$e^+ - e^-$ annihilation into
two photons
(511 keV)



Case 2

high energetic photons
1274.6keV
(Prompt)







TOT spectra for 511 keV and 1274.6 keV and their scattering angles



TOT spectra



Scattered angles (θ)



Sushil K. Sharma

- From the ^{22}Na source, one can get 511 keV from the $e^+ e^-$ annihilation and 1274.6 from prompt.
- TOT spectra resemble the Compton like structure, where *TOT* is the estimation of energy deposition.






TOT vs Energy deposition




Sushil K. Sharma





Positronium atom – a unique laboratory

- ❑ First time detected positronium in Gas : Martin Deutsch
Nobel prize in 1956 for discovering Ps
- ❑ Positronium is like hydrogen atom without nuclei consist of electron and positron.
- ❑ Eigenstate for C,P, CP operators
- ❑ Undergoes self-annihilation into gamma quanta








Tests of discrete symmetries with J-PET detector



Sushil K. Sharma





Selected recent publications -

P. Kowalski et al., *Phys. in Med. & Bio.* **63** (2018)

L. Raczyński et al., *Phys. Med. Bio.* **62** (2017) 5076

P. Moskal et al., *Phys. in Med. & Bio.* **61** (2016) 2025

A. Gajos et al., *Nucl. Inst. & Meth. In Phys. Res. A* **819**(2016) 54



G. Korcyl et al., *IEEE Trans. on Med. Imag.* (2018)

A. Wieczorek et al., *PLoS ONE* **12** (11): E0186728 (2017)


D. Kaminska et al., *Eup. Phys. J. C* **76** (2016)

J-PET's plastic revolution, Cern Courier, October 2018

Trilateration methods



Solving the GPS Geometry Problem



A. Gajos et al., Nucl. Inst. and Meth. A819 (2016) 54



Operators constructed using o-Ps spin



Sushil K. Sharma

Operators	C	P	T	CP	CPT
$\vec{S} \cdot \vec{k}_1$	+	-	+	-	-
$\vec{S} \cdot (\vec{k}_1 \times \vec{k}_2)$	+	+	-	+	-
$(\vec{S} \cdot \vec{k}_1) (\vec{S} \cdot (\vec{k}_1 \times \vec{k}_2))$	+	-	-	-	+




Example to test time reversal symmetry in o-Ps



Sushil K. Sharma

$$|\vec{k}_1\rangle > |\vec{k}_2\rangle > |\vec{k}_3\rangle$$





Additional operators with J-PET detector





Sushil K. Sharma

Operators	C	P	T	CP	CPT
$\vec{S} \cdot \vec{k}_1$	+	-	+	-	-
$\vec{S} \cdot (\vec{k}_1 \times \vec{k}_2)$	+	+	-	+	-
$(\vec{S} \cdot \vec{k}_1) (\vec{S} \cdot (\vec{k}_1 \times \vec{k}_2))$	+	-	-	-	+

Studies of discrete symmetries using the photon's polarization

Unique feature of the J-PET





Photon's Polarization

$$\vec{\epsilon} = \vec{k} \times \vec{k}'$$

P. Moskal et al., *Eur. Phys. J C* **78** (2018) 970

P. Moskal et al., *Acta. Phys. Polon. B* **47** (2016) 509, P. Moskal et al., arXiv : 1809.10397v1





Determining the Single photon's polarization

(Based on only the Compton scattering)




Sushil K. Sharma

- ✓ The scattering distribution of photons can be described by Klein-Nishina diff. cross section.
- ✓ The visibility to observe the angular correlation between scattering and polarization plane (φ) is a function of **Photon's energy** and its **scattering angle (θ)**.
- ✓ For **511 keV photon** and scattering **$\theta = 81.66^\circ$** , the Klein-Nishina differential xn for φ :





Eventwise observation of scattering angles of back-to-back photons





The positron emitting from the ^{22}Na source can annihilate into two photons, directly or through the formation of positronium atoms.

The interactions of the annihilated photons can be measured event-wise and allow to study the ***relative polarization b/w the photons.***



Relative polarization of entangled annihilated photons in decay of p-Ps atom



- Polarization vectors of annihilation photons are mutually orthogonal states.
- Photons mostly scatter at right angles to their electric field vector and this impose an **Expected angular correlation** between the scattering angles.
- With the J-PET detector we can measure scatterings of back-to-back photons and thus can study the angular correlation(φ) between the scattering angles/planes.
- Thus the **angle between two scattering planes(φ)** can be an estimator of relative polarization of two photons.



Experimental evidence for the measurement of photon's polarization with the J-PET detector





Small annihilation chamber was used






4 – hit events were studied





4 – hit events were studied




Assignment of 3rd HIT

Scatter test = $\text{time}_{\text{measured}} - \text{time}_{\text{calculated}}$



Assignment of 4th HIT



Relative angles between scattered planes *as a measure of* Relative polarization of annihilation photons



Courtesy to J. Raj
and N. Krawczyk





Eventwise observation of scattering angles of back-to-back photons



Sushil K. Sharma

Operators	C	P	T	CP	CPT
$\vec{S} \cdot \vec{k}_1$	+	-	+	-	-
$\vec{S} \cdot (\vec{k}_1 \times \vec{k}_2)$	+	+	-	+	-
$(\vec{S} \cdot \vec{k}_1) (\vec{S} \cdot (\vec{k}_1 \times \vec{k}_2))$	+	-	-	-	+


New operators available with J-PET

With


$$\vec{\epsilon}_i = \vec{k}_i \times \vec{k}'_i$$

$\vec{k}_2 \cdot \vec{\epsilon}_1$	+	-	-	-	+
$\vec{S} \cdot \vec{\epsilon}_1$	+	+	-	+	-
$\vec{S} \cdot (\vec{k}_2 \times \vec{\epsilon}_1)$	+	-	+	-	-

Porous material



Positronium life time spectra*





*K. Dulski et al., Hyperfine Interact 40 (2018) 239


Study of angular correlations among the photon originating from the decay of Positronium atom can provide an insight into the rare decays : e.g: p-Ps \rightarrow 3γ

Generated : D. Kaminska et al.,
Eur. Phys. J C 76 (2016) 445

In o-Ps decay : angular correlation



Experimental : Courtesy to J. Chhokar








Charge conjugation symmetry

Based on angular correlations and rate ratio $3\gamma / 2\gamma$



Study of angular correlations among the 3 photon originating from the decay of Positronium atom, distinguish based on the life time of positronium atom at various symmetrical configuration ($p\text{-ps} \rightarrow 3\gamma$ decay mode is restricted by Bose-statistics)

Positronium life time*



2 Decay Rate ratio : ?
 $3\gamma / 2\gamma$

* K. Dulski et al., Hyperfine Interact 40 (2018) 239






Charge conjugation symmetry


Based on angular correlations and rate ratio $3\gamma / 2\gamma$



Study of angular correlations among the 3 photon originating from the decay of Positronium atom, distinguish based on the life time of positronium atom at various symmetrical configuration ($p\text{-ps} \rightarrow 3\gamma$ decay mode is restricted by Bose-statistics)



Positronium life time*



* K. Dulski et al., Hyperfine Interact 40 (2018) 239






Modular J-PET – extention to 3 lyaer prototype




Sushil K. Sharma

24.11






Fourth Layer along with current prototype



**24 PORTABLE
modules**






Advantages with Modular J-PET

Geometry configuration made with Geant4 package




24 Modules placed as
the innermost layer



Advantage

- ✓ Several times better efficiency
- ✓ Standalone PET/detector

24 Modules rearranged
and add 2 layers



Multi-configurational



Summary



- ✓ A Positron Emission Tomograph based on plastic scintillators constructed and commissioned .
- ✓ Discrete symmetries are very crucial in order to understand the inequality between matter and anti-matter.
- ✓ Such inequality should have contribution of symmetry violation not only in baryonic and mesonic sectors but also from leptons.
- ✓ The experimental data on fundamental symmetry tests in leptonic sector is very scarce.
- ✓ The J-PET detector is capable to study the C, T, CP and CPT test in the decays of Ps atoms with better precision.
- ✓ Possibility to measure polarization direction will add up new scope to study the additional odd symmetric operators and phenomena like multi-particle entanglement.



J-PET collaboration



P. Moskal¹, M. Bala¹, C. Curceanu², E. Czerwiński¹, J. Chhokar, K. Dulski¹, A. Gajos¹, M. Gorgol³, B. Hiesmayr⁴, D. Kamińska¹, G. Korcyl¹, P. Kowalski⁵, T. Kozik¹, W. Krzemień⁵, E. Kubicz¹, M. Mohammed¹, N. Krawczyk¹, M. Pawlik-Niedźwiecka¹, Sz. Niedźwiecki¹, M. Pałka¹, L. Raczyński⁵, Z. Rudy¹, J. Raj¹, O. Rundel¹, N. Sharma¹, S. Sharma¹, Shivani¹, M. Silarski¹, J. Smyrski¹, A. Strzelecki¹, W. Wiślicki⁵, B. Zgardzińska³

¹Jagiellonian University, Poland; ²LNF INFN, Italy; ³Maria Curie-Skłodowska University, Poland;

⁴University of Vienna, Austria; ⁵National Centre for Nuclear Research, Poland;



Thank you for your attention